

AD 686472

A FORTRAN PROGRAM TO GENERATE TIME VS. SOLAR DEPRESSION TABLES

by
Allan R. Zeiner



NORTHEASTERN UNIVERSITY
Boston, Massachusetts 02115

Contract No. AF 19(628)-5731

Project No. 7661

Task No. 766102

Work Unit No. 76610201

SCIENTIFIC REPORT NO. 1

January 31, 1969

Distribution of this document is unlimited. It may be released to the
Clearinghouse, Department of Commerce, for sale to the general public.

Contract Monitor: Sam Silverman
Aeronomy Laboratory

Prepared
for

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES
OFFICE OF AEROSPACE RESEARCH
UNITED STATES AIR FORCE
BEDFORD, MASSACHUSETTS 01730

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va 22151

Qualified requestors may obtain additional copies from the Defense Documentation Center. All others should apply to the Clearinghouse for Federal Scientific and Technical Information.

ACCESSION No.	
CFSTI	WHITE SECTION <input checked="" type="checkbox"/>
DDC	GRAY SECTION <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY _____	
DISTRIBUTION/AVAILABILITY GROUP	
DIST.	AVAIL. CODE/NO. SPECIAL

AFCRL-69-0090

A FORTRAN PROGRAM TO GENERATE
TIME VS. SOLAR DEPRESSION TABLES

by
Allan R. Zeiner

NORTHEASTERN UNIVERSITY
Boston, Massachusetts 02115

Contract No. AF 19(628)-5731

Project No. 7661
Task No. 766102
Work Unit No. 76610201

SCIENTIFIC REPORT NO. 1

January 31, 1969

Distribution of this document is unlimited. It may be released to the
Clearinghouse, Department of Commerce, for sale to the general public.

Contract Monitor: Sam Silverman
Aeronomy Laboratory

Prepared
For

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES
OFFICE OF AEROSPACE RESEARCH
UNITED STATES AIR FORCE
BEDFORD, MASSACHUSETTS 01730

Abstract

A program is presented for the generation of tables of solar depression versus time on a particular day or period of days for any location. The input data required are two values for each day and the value of ΔT for the year as taken from "The American Ephemeris and Nautical Almanac," the longitude and latitude of the location of interest, the maximum and minimum zenith angle desired. The program is suitable for or easily adapted to any first generation or newer computer capable of being programmed in Fortran.

Table of Contents

Chapter	Page
I. Introduction - - - - -	1
II. Program Explanation - - - - -	1
III. Use of the Program - - - - -	2
IV. Appendix	
A. Program Flow Chart - - - - -	5
B. Program - - - - -	7
C. Sample Ephemeris Input Data - - - - -	13
D. Sample Output - - - - -	15
E. Sample Calculation - - - - -	17
F. Glossary of Terms - - - - -	18
Bibliography - - - - -	21

I. Introduction

The tabulated values of solar depression versus time currently available are of questionable value when used as a common denominator in the reduction of data gathered during atmospheric observations. In most cases, the discrete intervals of depression angle, day number and latitude are too widely spaced to allow for an accurate interpolation between given values. The accurate calculation of the time associated with a given depression angle, at a specific location, on a particular day or period of days has, up to now, been a laborious task. The program presented in this report has been developed to allow the generation of useful tables for any location. The input data required are two values for each day and the value of ΔT for the year as taken from "The American Ephemeris and Nautical Almanac" which is published in advance for any year, the longitude and latitude of the location of interest, the maximum and minimum zenith angles of interest and the increment desired between the given zenith angles.

II. Program Explanation

The basic method employed for computation is that illustrated in section 13.D of the "Explanatory Supplement to the Astronomical Ephemeris and Nautical Almanac." As can be seen upon checking the sample calculations, reproduced in Appendix E, the accuracy of this method is greatly influenced by the selection of the time used for the interpolation of the tabulated values of apparent declination and ephemeris transit. These tabulated values are taken from the Sun data for "0" hours ephemeris time as given in "The American Ephemeris and Nautical Almanac," hereafter abbreviated "AE and NA."

In order to achieve maximum accuracy, a reiterative method of calculation is used based upon the Newton Forward method of interpolation. An approximate time, used as an interpolation factor is inserted and with this, a calculated time for the zenith angle of interest is generated. This calculated time is then compared with the approximate time inserted, and unless the difference between the two is less than a specified tolerance, (0.0001 hours), the approximate time is replaced by the calculated time and the procedure repeated until the tolerance condition is met. Note that this leaves us with a value of ephemeris time for the angle of interest. The value of ΔT is now subtracted from the final calculated time to yield Greenwich mean time at the Greenwich meridian. To this value is added the value of West Longitude, converted to hours to finally yield the local Greenwich mean associated with the particular zenith angle. The program next increments to the next desired solar angle, and the procedure repeated until all desired angles for a given day are computed. Following this, the data for the next day are read in, and the calculations are repeated.

The zenith angle used to calculate sunrise and sunset is 90° , zero degrees horizon angle, rather than $90^{\circ} 50'$ as used in the Explanatory Supplement to the "AE and NA." This angle was chosen because the primary interest was in the center point of the sun's disk rather than the upper limb; therefore, the semi-diameter of $16'$ was neglected. The horizontal refraction of $34'$ was neglected because our interest was in the effect of the sun upon the upper atmosphere.

The horizon angles that are tabulated in the output are those for points at sea level. Since our interest was in zenith atmospheric measurements only, this suited our purpose. If, however, the altitude of the observing location is to be taken into consideration, the zenith maximum and minimum angles may be increased by a factor of $1.17h^{1/2}$, where h is measured in feet.

The basic formula used for computation is:

$$\cos(h) = -\tan(\delta) \tan(\varphi) + \cos(z) \cos(\delta) \sec(\varphi)$$

where φ is the latitude of interest,

δ is the apparent declination of the sun at the time of interest,

z is the desired angle of the sun as measured from zenith,

and h is the local hour angle of the sun.

In temperate latitudes, no check normally need be made of the value of h calculated. When dealing with latitudes greater than 60° , however, and when the zenith angles of interest are 90° to 108° , there are days of the year when these angles do not actually occur; that is, these zenith angles are not reached. In the generation of the tables, these angles are inserted in the basic formula as a value for (z) , resulting in a computed value of (h) which is also in error. Therefore, the time computed is also in error. This possibility has been considered, and the program will reject all erroneous times that are calculated due to the insertion of an angle that is not physically realizable, and insert a series of dash marks in the space ordinarily occupied by the time. See Appendix D for a sample output page which gives twilight information for Thule, Greenland, on September 10, 1968.

III. Use of the Program

This section is intended to provide all the information necessary to apply this program to any first generation or newer computer capable of being programmed in Fortran. However, the input and output statements used in the program as listed in Appendix B may have to be modified to allow its use on any specific computer. The body of the program may be used as listed.

Input data and format

The information given below is in the order that it necessarily must be placed

in a data deck. We will consider as an example the generation of a set of tables for the months of September and October, 1968, at a location of $58^{\circ} 40'$ latitude and $42^{\circ} 33'$ longitude. We desire the times associated with the horizon angles from -18° to 0° in $1/2^{\circ}$ increments. This would correspond to zenith angles of 108° to 90° .

Card 1. Longitude and latitude, with west longitude and north latitude being taken as positive.

Enter the data with input format (2F9,4). Example,

; : 42.5500; : 58.6667, where ":" indicates a space.

Card 2. Increment between tabulated angles, maximum zenith angle, minimum zenith angle, and value of ΔT as taken from the "AE and NA"

Enter the data with the input format (F4.1, 2F5.0, F9.4)

Example, :0.5:108; :90.;37.0000

Card 3. Month

Enter the month number, 1 thru 12, with the input format (I2).

Example, :9.

Cards 4⁺. Day of month, apparent declination in degrees, minutes, and seconds, and ephemeris transit in hours, minutes, and seconds.

Enter the data with the input format (I2, I5, I3, F5.1, 2I3, F6.2).

Example, :1 : : :18 : : 3:53.4:12 : : 6:13.83.

22	0 22 50.7 11 52 38.89
23	0-30-31.8 11 52 18.00
24	0-23-55.2 11 51 57.23
25	0-47-18.9 11 51 36.60
26	-1 10 42.7 11 51 16.14

34	-4 177 9.8 11 48 40.48
----	------------------------

50

10

1	-3 7 29.0 11 49 36.86
---	-----------------------

35	-15 1942.5 11 43 36.51
----	------------------------

99

Notice that there are four more days data than there are days in the month. These extra days are the first four days of the following month and are necessary for accurate interpolation of the input values of apparent declination and ephemeris transit.

Notice also the manner in which the sign of the apparent declination data changes. Unless the degrees term is an integer other than zero when the minutes and seconds are negative, the minutes and seconds terms must be given the minus sign.


Control

If tables are to be generated for several months during the same computer run, the last entry in the day number column for all months except the last must be 50, followed by the next month's data. This allows the following month's data to be fed in automatically. To stop the program at the end of the last month's data, enter a 99 in the day number column using the same format as the day number, (12). See Appendix C and the example given in the directions for cards 4⁺.


It should be obvious that there is no need to generate a whole month's table if the information is needed only for a small number of days. For example, if the period of interest were from Sept. 8 to Sept. 10, 1968, it would be necessary only to insert ephemeris data for Sept. 8 to Sept. 14, 1968. As explained previously, the extra four days data are used for the purposes of interpolation.

Appendix A

Program Flow Chart



Program Flow Chart (continued)



Appendix B

Program

```
DIMENSION IAD1(5), IAD2(5), AD3(5), APD(5), AD1(5), AD2(5)
DIMENSION IER1(5), IER2(5), ER3(5), ETR(5), ER1(5), ER2(5)
DIMENSION DAPD1(4), DAPD2(3), DAPD3(2), IDY(5)
DIMENSION DETR1(4), DETR2(3), DETR3(2)
c    THE LONGITUDE AND LATITUDE ARE NOW READ IN.
    READ 7, ALONG, ALAT
7    FORMAT (2F9.4)
c    THE ANGLE INCREMENT, ZENITH MAXIMUM AND MINIMUM
c    ANGLE AND VALUE OF DELTA T ARE NOW READ IN.
    READ 9, AINC, ZANGMX, ZANGMN, DELTAT
9    FORMAT (F4.1, 2F5.0, F9.4)
c    DEGREE TO RADIAN CONVERSION FACTORS AND INITIAL
c    VALUE OF APPROXIMATE TIME ARE COMPUTED.
    ZANGMX = ZANGMX + AINC
    DTOR = 3.1415926/180.
    RTOD = 180./3.1415926
    ATIME = 6.00
c    THE MONTH NUMBER AND THE FIRST FIVE DAYS
c    DATA ARE READ IN, TRIG. CONSTANTS COMPUTED,
c    AND OUTPUT HEADINGS PRINTED.
911  READ 912, MO
912  FORMAT(I2)
    DO 11 I = 1, 5
11   READ 10, IDY(I), IAD1(I), IAD2(I), AD3(I), IER1(I), IER2(I), ER3(I)
10   FORMAT (I2, I5, I3, F5.1, 2I3, F6.2)
    ALATRD = ALAT*DTOR
    A = (SINF (ALATRD))/(COSF (ALATRD))
    E = 1./COSF (ALATRD)
111  PRINT 890, ALONG, ALAT
890  FORMAT (1H1, 9HLOCATION, 5X, 9HLONGITUDE, F11.4, 5X,
1    8HLATITUDE, F11.4)
    PRINT 891
891  FORMAT (10X)
    PRINT 363
```

Program (continued)

```

363      FORMAT (1X, 5HMONTH, 2X, 3HDAY, 4X, 7HHORIZON, 5X,
1        7HMORNING, 13X, 7HEVENING)
      PRINT 365
365      FORMAT (15X, 5HANGLE, 7X, 11HHR MIN SEC, 9X, 11HHR MIN SEC)
      PRINT 366
366      FORMAT (/)
      ANG = 0.0
      SPACER = -1.0
c      ALL DATA IN MINUTES AND SECONDS ARE CONVERTED
c      TO FRACTIONS OF HOURS OR DEGREES AND A DIFFERENCE
c      TABLE IS COMPUTED.
12      DO 21 J = 1, 5
      AD1 (J) = IAD1 (J)
      AD2 (J) = IAD2 (J)
      ER1 (J) = IER1 (J)
      ER2 (J) = IER2 (J)
      IF (AD1(J) ) 18, 20, 20
18      APD (J) = AD1 (J) - (AD2 (J)/60.) - (AD3 (J)/3600.)
      GO TO 21
20      APD(J) = AD1(J) + (AD2(J)/60.) + (AD3(J)/3600.)
21      ETR(J) = ER1(J) + (ER2(J)/60.) + (ER3(J)/3600.)
      DO 26 K = 1, 4
      DAPD1 (K) = APD (K+1) - APD (K)
26      DETR1 (K) = ETR (K+1) - ETR (K)
      DO 30 L = 1, 3
      DAPD2(L) = DAPD1(L+1) - DAPD1(L)
30      DETR2(L) = DETR1(L+1) - DETR1(L)
      DO 34 M = 1, 2
      DAPD3(M) = DAPD2(M+1) - DAPD2(M)
34      DETR3(M) = DETR2(M+1) - DETR2(M)
      DAPD4 = DAPD3(2) - DAPD3(1)
      DETR4 = DETR3(2) - DETR3(1)
      D = ZAMGMZ*DTOR
c      AN INTERPOLATION FACTOR IS COMPUTED AND WITH
c      IT THE APPARENT DECLINATION AND EPHEMERIS TRANSIT
c      CORRESPONDING TO THE APPROXIMATE TIME ARE COMPUTED
38      B = ATIME/24. + 1.
      B1 = B-1.
      B2 = B-2.

```

Program (continued)

```

B3 = B-3.
B4 = B-4.
ITL = 0
ITLF = 0
AD = APD(1) + DAPD1(1)*B1 + DAPD2(1)*B1*B2/2.
1 +DAPD3(1)*B1*B2*B3/6. + DAPD4*B1*B2*B3*B4/24.
ET = ETR(1) + DETR1(1)*B1 + DETR2(1)*B1*B2/2.
1 +DETR3(1)*B1*B2*B3/6. + DETR4*B1*B2*B3*B4/24.
c USING THE VALUES OF APPARENT DECLINATION AND EPHEMERIS
c TRANSIT COMPUTED BY INTERPOLATION, THE LOCAL HOUR
c ANGLE IS COMPUTED AND CONVERTED TO AN EQUIVALENT TIME.
ADRAD = AD*DTOR
C = (SINF (ADRAD))/(COSF (ADRAD))
FE = 1./COSF (ADRAD)
G = COSF (D)
H COS = -A*C+E*FE*G
O = H COS**2
X = ABSF (1.-O)
HSIN = SQRTF (X)
ABHCOS = ABSF (H COS)
HRAD = ATANF (HTAN)
IF (H COS) 59, 61, 61
59 HDEGRE = 180. - HRAD*RTOD
GO TO 62
61 HDEGRE = HRAD*RTOD
62 HTIME = HDEGRE/15.
c IT IS NEXT DETERMINED WHETHER THE APPROXIMATE TIME
c USED CORRESPONDS TO A RISING OR SETTING ANGLE AND
c IF THE TOLERANCE CONDITIONS IS MET. IF NOT, THE
c APPROXIMATE TIME IS REPLACED BY THE COMPUTED TIME AND
c THE ENTIRE PROCESS REPEATED UNTIL THE TOLERANCE
c CONDITION IS MET.
IF(12. -ATIME) 70, 63, 63
63 TRISE = ET-HTIME
ABTIME = ABSF(TRISE - ATIME)
IF(.0001 - ABTIME) 66, 68, 68
66 ATIME = TRISE
GO TO 38
68 IF (ANG) 661, 661, 663

```


Program (continued)

```

661     ATIME1 = ATIME
663     ATIME = TRISE + 12.
        GO TO 38
70      TSET = ET + HTIME
        ACTIME = ABSF (TSET - ATIME)
        IF (.0001 - ACTIME) 73, 75, 75
73      ATIME = TSET
        GO TO 38
c       THE COMPUTED TIMES ARE CONVERTED TO GMT AND BROKEN
c       DOWN INTO HOURS, MINUTES AND SECONDS. NEXT A
c       CHECK IS MADE OF THE VALIDITY OF THE TIMES COMPUTED,
c       AN APPROPRIATE OUTPUT FORMAT STATEMENT SELECTED
c       AND THE TIMES PRINTED.
75      GMTRSE = TRISE = ALONG/15. - DELTAT/3600.
        GMTSET = TSET + ALONG/15. - DELTAT/3600.
        IF(24. - GMTSET) 744, 745, 745
744     GMTSET = GMTSET-24.
745     ITRISE = GMTRSE
        ATRISE = ITRISE
        BTRISE = GMTRSE - ATRISE
        BTRISE = BTRISE*60.
        IBTRSE = BTRISE
        BTRSE = IBTRISE
        CTRISE = (BTRISE - BTRSE)*60.
        ITSET = GMTSET
        ATSET = ITSET
        BTSET = GMTSET - ATSET
        BSET = BTSET*60.
        IBSET = BSET
        ABSET = IBSET
        CTSET = (BSET - ABSET)*60.
        IDAY = IDY(1)
        SPACER = SPACER + 1.0
        IF (2.0 - SPACER) 779, 779, 800
779     SPACER = 0.0
        PRINT 780
780     FORMAT (10X)
800     EDEP = (ZANGMX - ANG) - 90.
        ADEP = 90. -(ZANGMX - ANG)

```

Program (continued)

```

201 IF(ANG-(AINC/2.)) 200, 201, 201
202 IF(TRISE - TRISE1) 202, 203, 203
202 ITLE = 5
203 IF(TSET1 - TSET) 204, 205, 205
204 ITLF = 10
205 IF(ITLE + ITLF)806, 806, 807
806 PRINT 90, MO, IDAY, ADEP, ITRISE,
1 IBTRSE, CTRISE, ITSET, IBSET, CTSET
90 FORMAT (1X, 2(2X, I3) F8.1, 5X 2I4, F6.1, 6X, 2I4, F6.1)
GO TO 200
807 IF ((ITLE + ITLF) - 5) 200, 808, 810
808 PRINT 809, MO, IDAY, ADEP, ITSET, IBSET, CTSET
809 FORMAT (1X, 2(2X, I3), F8.1, 7X, 12H-- -- ----, 2I4, F6.1)
GO TO 200
810 IF ((ITLE + ITLF)-10) 200, 811, 813
811 PRINT 812, MO, IDAY, ADEP, ITRISE, IBTRSE, CTRISE
812 FORMAT (1X, 2(2X, I3), F8.1, 5X, 2I4, F6.1, 8X, 12H-- -- ----)
GOTO 200
813 IF ((ITLE + ITLF)-15) 200, 814, 814
814 PRINT 815, MO, IDAY, ADEP
815 FORMAT (1X, 2(2X, I3), F8.1, 7X, 12H-- -- ----,
1 8X, 12H-- -- ----)
c NEXT IT IS DETERMINED IF ALL THE ANGLES FOR THE DAY
c HAVE BEEN COMPUTED. IF NOT, THE PROGRAM INCREMENTS
c TO THE NEXT ONE AND THE TIMES ARE COMPUTED.
c IF ALL THE ANGLES FOR THE DAY HAVE BEEN COMPUTED,
c THE PROGRAM INCREMENTS TO THE NEXT DAYS DATA
c AND CONTINUES AS BEFORE.
200 TRISE1 = TRISE
TSET1 = TSET
91 IF (ZANGMN - (EDEP + 90.))92, 95, 95
92 D = D-AINC*DTOR
930 ANG = ANG + AINC
ATIME = TRISE
GO TO 38
95 ATIME = TRISE
DO 102 J 1, 4
IDY(J) = IDY(J+1)
IAD1(J) = IAD1 (J+1)

```

Program (continued)

```
IAD2(J) = IAD2(J+1)
AD3(J) = AD3(J+1)
IER1(J) = IER1(J+1)
IER2(J) = IER2(J+1)
102  ER3(J) = ER3(J+1)
      READ 10, IDY(5), IAD1(5), IAD2(5), AD3(5 ), IER1(5), IER2(5), ER3(5)
      IF (IDY(5) - 50) 104, 125, 126
104   SPACER = -1.0
      GO TO 111
125   GO TO 911
126   STOP
      END
```

Appendix C

Sample Ephemeris Input Data

9
1 8 20 35.3 11 59 54.60
2 7 58 46.8 11 59 35.40
3 7 36 50.7 11 59 15.92
4 7 14 47.4 11 58 56.17
5 6 52 37.1 11 58 36.17
6 6 30 20.2 11 58 15.94
7 6 7 56.9 11 57 55.51
8 5 45 27.6 11 57 34.90
9 5 22 52.4 11 57 14.12
10 5 0 11.8 11 56 53.21
11 4 37 25.9 11 56 32.18
12 4 14 35.2 11 56 11.05
13 3 51 39.9 11 55 49.85
14 3 28 40.3 11 55 28.60
15 3 5 36.8 11 55 7.30
16 2 42 29.7 11 54 46.00
17 2 19 19.3 11 54 24.69
18 1 56 6.0 11 54 3.41
19 1 32 50.2 11 53 42.18
20 1 9 32.1 11 53 21.00
21 0 46 12.2 11 52 59.90
22 0 22 50.7 11 52 38.89
23 0 0-31.8 11 52 18.00
24 0-23-55.2 11 51 57.23
25 0-47-18.9 11 51 36.60
26 -1 10 42.7 11 51 16.14
27 -1 34 6.1 11 50 55.85
28 -1 57 28.8 11 50 35.76
29 -2 20 50.5 11 50 15.88
30 -2 44 10.6 11 49 56.24
31 -3 7 29.0 11 49 36.86
32 -3 30 45.2 11 49 17.76
33 -3 53 58.9 11 48 58.96
34 -4 17 9.8 11 48 40.48
50

Sample Ephemeris Input Data (continued)

10

1	-3	7	29.0	11	49	36.86
2	-3	30	45.2	11	49	17.76
3	-3	53	58.9	11	48	58.96
4	-4	17	9.8	11	48	40.48
5	-4	40	17.5	11	48	22.35
6	-5	3	21.8	11	48	4.60
7	-5	26	22.2	11	47	47.24
8	-5	49	18.6	11	47	30.29
9	-6	12	10.4	11	47	13.79
10	-6	34	57.5	11	46	57.75
11	-6	57	39.4	11	46	42.20
12	-7	20	15.8	11	46	27.14
13	-7	42	46.3	11	46	12.62
14	-8	5	10.5	11	45	58.63
15	-8	27	28.1	11	45	45.21
16	-8	49	38.7	11	45	32.37
17	-9	11	41.9	11	45	20.13
18	-9	33	37.2	11	45	9.50
19	-9	55	24.4	11	44	57.50
20	-10	17	2.9	11	44	47.14
21	-10	38	32.4	11	44	37.45
22	-10	59	52.4	11	44	28.41
23	-11	21	2.6	11	44	20.06
24	-11	42	2.4	11	44	12.39
25	-12	2	51.6	11	44	5.43
26	-12	23	29.6	11	43	59.17
27	-12	43	56.1	11	43	53.62
28	-13	4	10.5	11	43	48.81
29	-13	24	12.6	11	43	44.73
30	-13	44	1.8	11	43	41.41
31	-14	3	37.9	11	43	38.85
32	-14	23	0.3	11	43	37.07
33	-14	42	8.8	11	43	36.08
34	-15	1	3.0	11	43	35.89
35	-15	19	42.5	11	43	36.51

99

Appendix D

Sample Output

Location,		Longitude	68.6667			Latitude	76.5500		
Month	Day	Horizon Angle	Morning				Evening		
			Hr	Min	Sec		Hr	Min	Sec
9	10	-18.0	--	--	---		--	--	---
9	10	-17.5	--	--	---		--	--	---
9	10	-17.0	--	--	---		--	--	---
9	10	-16.5	--	--	---		--	--	---
9	10	-16.0	--	--	---		--	--	---
9	10	-15.5	--	--	---		--	--	---
9	10	-15.0	--	--	---		--	--	---
9	10	-14.5	--	--	---		--	--	---
9	10	-14.0	--	--	---		--	--	---
9	10	-13.5	--	--	---		--	--	---
9	10	-13.0	--	--	---		--	--	---
9	10	-12.5	--	--	---		--	--	---
9	10	-12.0	--	--	---		--	--	---
9	10	-11.5	--	--	---		--	--	---
9	10	-11.0	--	--	---		--	--	---
9	10	-10.5	--	--	---		--	--	---
9	10	-10.0	--	--	---		--	--	---
9	10	-9.5	--	--	---		--	--	---
9	10	-9.0	--	--	---		--	--	---
9	10	-8.5	--	--	---		3	41	43.0
9	10	-8.0	5	31	50.9		3	11	30.1
9	10	-7.5	5	59	12.1		2	49	34.6
9	10	-7.0	6	20	3.4		2	31	20.3
9	10	-6.5	6	37	42.6		2	15	17.9
9	10	-6.0	6	53	22.8		2	0	45.1
9	10	-5.5	7	7	40.2		1	47	17.8
9	10	-5.0	7	20	56.4		1	34	40.9

Sample Output (continued)

Location		Longitude	68.6667			Latitude	76.5500		
Month	Day	Horizon Angle	Morning				Evening		
			Hr	Min	Sec	Hr	Min	Sec	
9	10	-4.5	7	33	25.2	1	22	43.9	
9	10	-4.0	7	45	16.0	1	11	19.4	
9	10	-3.5	7	56	35.9	1	0	21.9	
9	10	-3.0	8	7	30.0	0	49	47.1	
9	10	-2.5	8	18	2.3	0	39	31.5	
9	10	-2.0	8	28	16.1	0	29	32.4	
9	10	-1.5	8	38	14.2	0	19	47.5	
9	10	-1.0	8	47	58.5	0	10	15.0	
9	10	-0.5	8	57	31.1	0	0	53.1	
9	10	- 0	9	6	53.5	23	51	40.4	

Appendix E

Sample Calculation

Reproduction of example from section 13.D of the Explanatory Supplement to the American Ephemeris and Nautical Almanac.

	March 7, 1960		Latitude 52°	
$\phi = 52^{\circ}$	$\tan \phi = 1.2799$		$\sec \phi = 1.6243$	
	Morning Twilight	Sunrise	Sunset	Evening Twilight
Approx. time	4 ^h 30 ^m	6 ^h 30 ^m	18 ^h 00 ^m	19 ^h 30 ^m
δ (A.E., pg 21)	-5 ^o 17'	-5 ^o 15'	-5 ^o 04'	-5 ^o 02'
tangent δ	-0.0925	-0.0919	-0.0887	-0.0881
secant δ	1.0043	1.0042	1.0039	1.0039
z	108 ^o	90 ^o 50'	90 ^o 50'	108 ^o
cosine z	-0.3020	-0.0145	-0.0145	-0.3090
$-\tan \phi \tan \delta$	0.1184	0.1176	0.1135	0.1128
$\cos z \sec \phi \sec \delta$	-0.5041	-0.0237	-0.0237	-0.5039
cosine h	-0.3857	-0.0939	0.0898	-0.3911
	h min	h min	h min	h min
h	7 30.7	5 38.5	5 39.4	7 32.1
ephemeris transit	12 11.2	12 11.2	12 11.0	12 11.0
Time (local)	4 40.5	6 32.7	17 50.4	19 43.1

Appendix F

Glossary of Terms

IAD1	apparent declination, degrees (fixed point)
IAD2	apparent declination, minutes (fixed point)
AD3	apparent declination, seconds
APD	apparent declination, degrees and decimals of degrees
AD1	apparent declination , degrees (floating point)
AD2	apparent declination, minutes (floating point)
IER1	ephemeris transit, hours (fixed point)
IER2	ephemeris transit, minutes (fixed point)
ER3	ephemeris transit, seconds
ETR	ephemeris transit, hours and decimals of hours
ER1	ephemeris transit, hours (floating point)
ER2	ephemeris transit, minutes (floating point)
DAPD1	apparent declination, first difference
DAPD2	apparent declination, second difference
DAPD3	apparent declination, third difference
DAPD4	apparent declination, fourth difference
DETR1	ephemeris transit, first difference
DETR2	ephemeris transit, second difference
DETR3	ephemeris transit, third difference
DETR4	ephemeris transit, fourth difference
IDY	day number
ALONG	longitude
ALAT	latitude
AINC	degree increment
ZANGMX	maximum zenith angle
ZANGMN	minimum zenith angle
DELTAT	correction factor, ephemeris time to universal time
DTOR	conversion factor, degrees to radians
RTOD	conversion factor, radians to degrees
ATIME	approximate time
MO	month
ALATRD	latitude in radians
A	tangent (latitude)
E	secant (latitude)
ANG	counter for angle increment

Glossary of Terms (continued)

SPACER	counter to control output spacing
D	zenith angle in radians
B	interpolation factor
B1 TO B4	interpolation factors
ITLE	error check counter
ITLF	error check counter
AD	apparent declination after interpolation
ET	ephemeris transit after interpolation
ADRAD	apparent declination in radians
C	tangent (apparent declination)
FE	secant (apparent declination)
G	cosine (zenith angle)
HCOS	cosine of hour angle
HSIN	sine of hour angle
ABHCOS	absolute value of the cosine of the hour angle
HTAN	tangent of hour angle
HRAD	hour angle in radians
HDEGRE	hour angle in degrees
HTIME	time corresponding to hour angle of H
TRISE	local time of zenith angle during AM
ABTIME	absolute value of local time during AM
TSET	local time of zenith angle during PM
ACTIME	absolute value of local time during PM
GMTRSE	GMT of hour angle during AM
GMTSET	GMT of hour angle during PM
ITRISE	hour of AM zenith angle (fixed point)
ATRISE	hour of AM zenith angle (floating point)
BTRISE	minutes of AM zenith angle (floating point)
IBTRSE	minutes of AM zenith angle (fixed point)
BTRSE	minutes of AM zenith angle (floating point)
CTRISE	seconds of AM zenith angle (fixed point)
ITSET	hours of PM zenith angle (fixed point)
ATSET	hours of PM zenith angle (floating point)
BTSET	minutes of PM zenith angle (floating point)
IBSET	minutes of PM zenith angle (fixed point)
BSET	minutes of PM zenith angle (floating point)
ABSET	minutes of PM zenith angle (floating point)
CTSET	seconds of PM zenith angle (floating point)

Glossary of Terms (continued)

IDAY	day for printout
EDEP	counter for zenith angle
ADEP	counter for zenith angle
TRISE1	ne of zenith angle AM
TSET1	time of zenith angle PM

Bibliography

Her Majesty's Stationary Office. Explanatory Supplement to the Astronomical Ephemeris and the American Ephemeris and Nautical Almanac. Prepared jointly by the Nautical Offices of the United Kingdom and the United States of America. London: Her Majesty's Stationary Office, 1961.

U.S. Government Printing Office. The American Ephemeris and Nautical Almanac for the Year 1968. Prepared jointly by the Nautical Almanac Offices of the United Kingdom and the United States of America. Washington: Government Printing Office, 1966.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Northeastern University Boston, Massachusetts 02115		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE A FORTRAN PROGRAM TO GENERATE TIME VS. SOLAR DEPRESSION TABLES		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Scientific. Interim.		
5. AUTHOR(S) (Last name, first name, initial) Allan R. Zeiner		
6. REPORT DATE 31 January 1969	7a. TOTAL NO. OF PAGES 23	7b. NO. OF REFS 2
8a. CONTRACT OR GRANT NO. AF19(628)-5731 a. Project, Task, Work Unit Nos. 7661-02-01 c. DoD Element 6240539F d. DoD Subelement 681000		9a. ORIGINATOR'S REPORT NUMBER(S) Scientific Report No. 1
		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) AFCRL-69-0090
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.		
11. SUPPLEMENTARY NOTES TECH, OTHER		12. SPONSORING MILITARY ACTIVITY Air Force Cambridge Research Laboratories (CRA) L. G. Hanscom Field Bedford, Massachusetts 01730
13. ABSTRACT A program is presented for the generation of tables of solar depression versus time on a particular day or period of days for any location. The input data required are two values for each day and the value of ΔT for the year as taken from "The American Ephemeris and Nautical Almanac," the longitude and latitude of the location of interest, the maximum and minimum zenith angle desired. The program is suitable for or easily adapted to any first generation or newer computer capable of being programmed in Fortran.		

DD FORM 1473
1 JAN 64

Unclassified

Security Classification

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Solar Depression Angles Solar Depression Times Fortran Program						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

Unclassified

Security Classification